

Co-varying eye movements and power modulations of alpha oscillations during working memory: a pilot study

Arne D. Hansen¹, Dawid Strzelczyk^{1,2}, Lea Z. M. Bächlin¹, Nicolas Langer^{1,2} & Tzvetan Popov^{1,2}

Contact

Arne Hansen
arne.hansen2@uzh.ch

1: Methods of Plasticity Research, Department of Psychology, University of Zurich, Zurich, Switzerland
2: Neuroscience Center Zurich (ZENZ), Zurich, Switzerland

1. Introduction

- There is an **inconsistency** in the literature regarding the **direction of the association between alpha power and WM load**.^[1]
- The present proposal suggests that **different gaze patterns during commonly used WM tasks** (i.e., Sternberg and N-back) **predict variations in alpha power**.
- Differences in alpha power with WM varied with **variability in oculomotor action**: the higher the variability the stronger the decrease in alpha power and vice versa.^[2]
- The hypothesis of this study consequently is that variation of alpha power with WM load, or more generally, power modulations appear correlated with the cognitive task at hand, while in fact they evolved **primarily to support oculomotor control**.




Fig. 1: Increases and decreases in alpha amplitude during working memory retention, depending on WM load and recruited sensory area, showcasing direction of the association between alpha power and WM load. Adapted from van Ede, 2018.

2. Hypotheses




Fig. 2: Sternberg and N-back show similar behavioral data but the direction of the association between alpha power and WM load differs between the tasks. It is hypothesized that these differences are due to distinct variability of oculomotor action.

3. Methods

Dataset

- N = 10 pilot participants (mean age = 25.9, ± 2.13, 50% female)

Sternberg Task




Fig. 3: Sternberg task. A trial consisted of a sequential presentation of 1, 4 or 7 letters (1.2 s each). The letters and the load were randomized. After a retention interval of 3 s another letter was presented. Participants were instructed to indicate by button press whether they thought it was a match or not.

N-back Task




Fig. 4: N-back task. A block consisted of the sequential presentation of 100 letters (2 s each). Depending on the condition (1-back, 2-back and 3-back), participants were instructed to indicate by button press whether they thought the currently presented letter was a match to the letter N-back.

Psychophysiological Data

- 128-channel ANT Neuro EEG system, sampling rate 500 Hz
- Eye tracker EyeLink 1000 Plus
- Preprocessing: BP FIR filter 1-45 Hz, Automagic^[3]

Behavioral Data

- Reaction time
- Accuracy

4. Results

Sternberg Reaction Times and Accuracy




Fig. 5: Sternberg reaction time in ms and accuracy values in correct percentage for WM load conditions 1 and 7.

Sternberg Grand Average




Fig. 7: Grand average occipital power spectra and topographical alpha power distribution across WM load conditions 1 and 7.

Sternberg Single Subjects




Fig. 9: Individual power spectra as function of WM load during the Sternberg task.

Gaze density during Sternberg task




Fig. 11: A | Sternberg WM load 1 gaze density. B | Sternberg WM load 7 gaze density. C | Absolute gaze density difference from WM load 7 (B) – WM load 1 (A). D | T-values of gaze density difference from WM load 7 (B) – WM load 1 (A).

N-back Reaction Times and Accuracy




Fig. 6: N-back reaction time in ms and accuracy values in correct percentage for 1-back and 3-back conditions.

N-back Grand Average




Fig. 8: Grand average occipital power spectra and topographical alpha power distribution across 1-back and 3-back conditions.

N-back Single Subjects




Fig. 10: Individual power spectra as function of WM load during the N-back task.

Gaze density during N-back task




Fig. 12: A | 1-back gaze density. B | 3-back gaze density. C | Absolute gaze density difference from 3-back (B) – 1-back (A). D | T-values of gaze density difference from 3-back (B) – 1-back (A).

References

- van Ede, F. (2018). Mnemonic and attentional roles for states of attenuated alpha oscillations in perceptual working memory: A review. *European Journal of Neuroscience*, 48(7), 2509–2515.
- Popov, T., Miller, G. A., Rockstroh, B., Jensen, O., & Langer, N. (2021). Alpha oscillations link action to cognition: An oculomotor account of the brain's dominant rhythm. *Neuroscience*.
- Pedroni, A., Bahreini, A., and Langer, N. (2019). Automagic: Standardized preprocessing of big EEG data. *NeuroImage*, 200:460–473.

5. Conclusion (preliminary)

- Gaze and alpha power modulation during WM covary
- High WM load is associated with stronger alpha power reduction and higher gaze variability in the N-back task
- High WM load is associated with lower gaze variability in the Sternberg task; significant increases in alpha power could not yet be shown
- Strong inter-individual variability with modulation of posterior alpha power with WM load